



CASUISTIC PAPER

Nathália Aparecida Ileck ^{1(ABCDG)}, Marcus Vinícius Gonçalves Torres de Azevedo ^{1(ABCDG)},
Dérick Patrick Artioli ^{1(ABCDG)}, Gladson Ricardo Flor Bertolini ^{2(ADFG)}

Lian Gong for treatment of fibromyalgia – a case study

¹ Centro Universitário Lusíada (Unilus), Santos, São Paulo, Brazil

² Universidade Estadual do Oeste do Paraná (Unioeste), Cascavel, Paraná, Brazil

ABSTRACT

Introduction. As a Complementary Integrative Practices (CIPs), Lian Gong has been increasingly used in the practice of Physiotherapy.

Aim. This study aims to verify the effects of Lian Gong in a patient with fibromyalgia.

Description of the case. In this case study the patient (one patient is evaluated) was diagnosed with fibromyalgia 29 years ago and sought care following discontent with previous treatments. Pain intensity was assessed with a visual analog scale (VAS), quality of life with SF-36, and the pressure pain threshold algometer (PPT). There were 16 visits with two weekly 60-minute sessions. When the patient was reevaluated, a Global Perception of Change (GPC) scale was added to assess general health. A folder was submitted to the patient for follow-up of exercises at home and asked to return after 4 weeks (follow-up period).

Conclusion. The results indicate improvement in pain, functional capacity and general health. As a first therapy treatment, Lian Gong proved promising results in one case of fibromyalgia. The possible benefits when combined with other forms of care should be explored by clinical trials to expand knowledge of health benefit potential.

Keywords. fibromyalgia, modalities of physiotherapy, Qigong, quality of life, rehabilitation, traditional chinese medicine

Introduction

Fibromyalgia is a chronic, rheumatic disease of undefined cause, likely to occur due to abnormalities in neuroendocrine regulation and as a response to stress, and has a higher prevalence in women between 40 and 60 years of age.¹ Despite the rheumatology approach, there are common neuropathic symptoms (paresthesia, numbness, and thermal sensitivity). Its incidence is 2-4% in the general population, with diffuse muscle pain as its main characteristic.² Genetic pre-disposition, emotional

and cognitive factors, body-mind relationship, and ability to cope with stressful situations are all characteristics that contribute to the onset of fibromyalgia, but its etiology remains uncertain.³

Besides the above, there are other symptoms such as fatigue, sleep disorders, depression, changes such as hyperalgia, allodynia, morning stiffness, and headache (which can be exacerbated by hormones), physical and mental stress, temperature changes, changes in diet and sleep window.¹ The diagnosis is based on a set of symp-

Corresponding author: Gladson Ricardo Flor Bertolini, e-mail: gladson_ricardo@yahoo.com.br

Participation of co-authors: A – Author of the concept and objectives of paper; B – collection of data; C – implementation of research; D – elaborate, analysis and interpretation of data; E – statistical analysis; F – preparation of a manuscript; G – working out the literature; H – obtaining funds

Received: 13.04.2021 | Accepted: 28.04.2021

Publication date: June 2021




Fig. 1A. Booklet with representation of the exercises to be performed by the participant

toms that need to be present for at least three months, with pain in both hemicorps, tender points in eleven (or more) specific areas, generalized fatigue, depression and anxiety.³

As there is no cure for fibromyalgia, the treatment is based on symptom control and improves quality of life with drug management and non-pharmacological therapies. Physical exercise is part of the forms of treatment indicated in the multidisciplinary approach, as well as low power laser, balneotherapy and cognitive-behavioral therapy. Besides the usual practices of aerobic activity, strengthening and stretching, alternative and complementary therapies (Acupuncture, Massage, Meditation, Tai Chi Chuan, Shiatsu, among others) are applied in the management of fibromyalgia. A recent literature review on the use of such therapies in fibromyalgia has pointed to promising results, however, with the exception that these effects were based on small studies, but given the low risk, they could be considered an adjunct form of treatment. Although Lian Gong

was briefly mentioned, there were no major considerations on the subject.¹ In recent years, there has been considerable growth in the use of resources beyond conventional Western medicine in rheumatic patients, yet scientific evidence of these practices remains under investigation. Lian Gong is considered one of these mind-body interventions that are part of the list of alternative treatments possibilities, consisting of gentle movements, breathing exercises, and meditation, producing gains in muscular power and endurance, flexibility, balance, and affecting psychological factors such as distress and anxiety.^{1,5-8}

Lian Gong is of Chinese origin, has a long history (more than five thousand Years), and is practiced by more than 60 million people in China. Lian Gong can be defined as the ability to release, strengthen, and direct Qi “life energy” through specific exercises, harmonizing breathing, posture, body movements, and mind. Qi flows through meridians and its balance determines good overall health according to Traditional Chinese



Fig. 1B. Booklet with representation of the exercises to be performed by the participant

Medicine (TCM).⁸⁻¹⁰ With this line of reasoning, the technique would be in a position to accelerate the self-healing process⁶. They are simple, smooth movements, with a choreographed, rhythmic routine, with respiratory control and can be adapted depending on the needs of the practitioner.^{6,7} Although its effectiveness continues to be investigated, it is aimed at improving physical and mental well-being and quality of life.⁸

In Brazil, one of the strategies incorporated for health promotion by the National Policy of Integrative and Complementary Practices (NPICP), better known as “CIPs”, published in the form of Ministerial Ordinances No. 971 on May 3, 2006 and No. 1,600 on July 17, 2006.⁹ Some Brazilian cities have implemented, within their health policies, the use of Lian Gong for the population.^{10,11}

While the effects of Lian gong in the West are still being explored, this resource is indicated in cases of fibromyalgia.⁸

Aim

The objective of this study was to present the effects of the technique, as a single therapy, in a patient with fibromyalgia that had been ongoing for decades.

Description of the case

The present study is characterized as a case study, being previously approved by the Research Ethics Committee of Centro Universitário Lusíada – UNILUS under number 2,243,010. The patient chosen was on the waiting list of the UNILUS Physiotherapy Clinic, with a diagnosis of fibromyalgia.

The patient E.M.L, 46 years, with 1.63 m height and 67 kg, had been diagnosed at 17 years of age, and had performed physiotherapeutic treatment (infrared, TENS and hydrotherapy) and medications (muscle relaxant and antidepressant) before, but remained dissatisfied with the results.

Before the evaluations began, the procedures were explained to the patient and the informed consent form




Fig. 1C. Booklet with representation of the exercises to be performed by the participant

was signed. We applied the visual analog pain scale (VAS - ranging from 0 (no pain) to 10 (maximum pain possible), SF-36 quality of life questionnaire (0 - 100, which evaluates functional capacity, physical aspects, pain, general health status, vitality, social, emotional and mental health) and the pressure pain threshold (PPT) (Tester® - Kgf) on the 18 points related to the tendon points of individuals with fibromyalgia (adding the maximum pressure supported at all points, to generate a single final value).^{5,9,16}

We also analyzed which Lian Gong movements the patient would be able to perform correctly or need only minimal adjustments, and then the treatment plan was constituted. After the second meeting, the sessions lasted 60 minutes, performing exercises that included the axial and appendicular skeleton, respecting the painful limit. It is therefore recommended that the session be made of six exercises in three series, totaling 18 exercises. For the evaluation of the exercises applicable to the patient in question, each movement was harmonious-

ly performed three times, with respiratory control and following the work sequence of the “anterior and posterior part” of Lian gong’s manual, totaling about 20 exercises per session.¹¹ The treatment was performed for eight weeks, twice a week, reaching 16 sessions.

In the reevaluation, in addition to the items mentioned above, the Global Perception of Change (GPC) scale was added (ranging from -5 to + 5, positive values indicate greater satisfaction), helping to analyze how efficient the treatment was.¹³ After the reevaluation and the end of the clinic visits, the patient received a booklet with photos (figure 1A–C) showing how to perform the exercises, which she was already familiar with, so that she could continue at home and return for a new reevaluation after four weeks (follow-up). She was also guided to access the following links in case of doubt in the execution: <https://www.youtube.com/watch?v=FgQkY-SAmJ6o> <https://www.youtube.com/watch?v=rbXOUi-VFLDo>.

The other instrument used for general pain assessment, VAS, showed a significant pain reduction effect, occurring 50.9% and 58.4% of reduction at completion and follow-up after treatment, respectively. The SF-36 questionnaire recorded increases (Functional Capacity and General Health Status), stabilization (Physical, Social, Emotional and Mental Health Aspects) and decrease (Vitality) of its values (Table 1).

Table 1. Quality of life assessment through SF-36. Higher scores indicate improved quality of life

	Begin	End	Follow-up
Functional capacity	80	80	90
Physical aspects	100	100	100
Improvement of pain	51	51	51
General health status	30	47	57
Vitality	95	90	90
Social aspects	100	100	100
Emotional aspects	100	100	100
Mental health	84	92	84

The evaluation of the pain pressure threshold recorded a progressive increase in the capacity to withstand a pressure of 21.6% at the end of clinic visits and a total of 23.1% when returning for follow-up evaluation. The improvement in pain accompanied by VAS and PPT, probably participation in the positive response, was noted and maintained at the end and the return to the last evaluation of the study, by grading its improvement by +3 on the GPC scale (Table 2).

Table 2. Parameters linked to pain and general improvement*

	Begin	End	Follow-up
VAS	5.3	2.6	2.2
PPT	1315	1600	1620
GPC	*	+3	+3

*VAS – visual analogue scale, PPT – pressure pain threshold, GPC – global pain change

Discussion

Use of Lian Gong has proven useful as an add on therapy in the treatment of fibromyalgia in the case studied. These findings corroborate those found in other studies, which describe positive results with Lian Gong among the possibilities of alternative and complementary therapies when treating fibromyalgia.¹⁴ The fact that it does not prescribe any specific device or tool for its performance, makes its practice valid in any environment and time. Through TCM, Lian Gong would be able to elevate the body's physical energy by promoting the fluidity of Qi and blood circulation, as showed by the increased electrical conductivity of meridians. It has already proven effective in preventing bone mineral loss, reducing

oxidative stress, increasing antioxidant enzymes, homeostasis of the autonomic nervous system, activation of immune system cells, such as in reducing pain, improving balance, flexibility, agility, strength, fatigue, quality of life and sleep.^{4,5}

This potentially justifies the improvements in overall pain, functional capacity, and general health status of the SF-36. Generalized chronic pain is the main complaint of fibromyalgia patients and is related to decreased quality of life. This association has previously been demonstrated in the comparison between the Fibromyalgia Impact Questionnaire (FIQ) and the VAS.¹⁵ Despite the short intervention period (8 weeks) and follow-up (4 weeks), these were already sufficient for pain to be minimized at half of the initial assessment, corroborated by improvement in the other instruments (PPT and GPC). Since with the severity of the pain, there is a reduction in functional capacity. The painful improvement is assumed to have contributed to the increase in Functional Capacity in the SF-36.

The SF-36 was lower than the initial value, only in the item Vitality, there was improvement in pain measured both by the VAS and the pressure gauge and satisfaction with the proposed treatment via the Global Perception of Change scale. This way, it is believed that this variable did not present much of a change because the data from the other domains initially already presented high values. Since diffuse pain is usually the main complaint of these patients and although it has not been altered via the SF-36, the benefit achieved by other assessment instruments (VAS, PPT and GPC) stands out. This improvement in pain complaint, added to the increase in Functional Capacity and the General Health Status, generates a promising idea of the effects of Lian Gong on fibromyalgia.

Commonly used in association with VAS, the pressure algometer in the comparison of individuals with fibromyalgia to rheumatoid arthritis, dyspareunia and controls, shows lower pressure tolerance values, i.e., they feel more pain with smaller stimuli.² The characterization of fibromyalgia is generalized pain in 4 out of 5 regions (4 quadrants and axial).²⁰ The increase in pain pressure threshold with the practice of Lian Gong generates optimism and the need for studies with a larger number of participants and time, to determine how much improvement is expected to the longevity of a mind-body exercise routine.

GPC is easily and quickly applied, described as clinically relevant in cases of fibromyalgia when related to the patient's general clinical picture and previously used with EVA and body map (18 common tender point sites). The higher the values obtained through the GPC, the better the pain (via VAS and painful points) and all other measures of fibromyalgia severity.¹³ What was found was in fact improvement of the painful condition (VAS and PPT) and the GPC.

Since fibromyalgia is not only musculoskeletal issue, there is psychoemotional involvement, Lian Gong fits the demand to be a mind-body therapy.⁴ Mental Health of the SF-36 showed this effect during the face-to-face period of the intervention, but it returned to initial values during the four weeks of follow-up. Therefore, the result may have been the engagement of the professional during the sessions and not restricted to the technique. It is noteworthy that there are indications for the exercise to be performed regularly and even daily, in the present study the indicated practice was regular, but with 2 sessions per week.⁸

Since fibromyalgia is not a curable disease, therapies that help control symptoms and improve quality of life are the focus for these patients.¹ The benefits gained from the practice of Lian Gong, its adaptability, and low cost, have caused this alternative to spread and conquer followers in Brazil.¹⁰ CIPs such as Lian Gong, act in the promotion and integrality of health, surpassing the rooted curative model as the assertive one. It is common for Lian Gong to be practiced in groups, providing socialization, strengthening of the professional-patient relationship, and amplification of the individual's notion of his or her general state of health through contact with others in similar conditions. The use of Lian Gong in cases of fibromyalgia strengthens the understanding of the need for active participation in the care process adding to medical approaches. Although the publication of the National Policy of Integrative and Complementary Practices in SUS was made in 2006 and Lian Gong is not a new therapy, both the training of professionals focused on these practices and the scientific basis require investment.

The methodology used for monitoring the case reported here, while effective in reflecting symptomatic changes, could have been more complete if it had used specific instruments that evaluated sleep, fatigue, anguish, anxiety, and depression, which are to the painful complaint and the SF-36 items are to be addressed.^{1,4,5} In addition to this limitation, it should be taken into consideration that this is a case study and cannot extrapolate the findings to all patients with fibromyalgia. Clinical trials will be more reliable in this response and it is worth noting the need for a control group to evaluate the real effects of this therapy, both its evaluation as an isolated form of treatment, as proposed here, although it is commonly described as adjunct. Progress has been made in proposing a placebo form of Lian Gong.⁴

Conclusion

Lian Gong stands out for its adaptability, easy application, low demand for the necessary structure, low cost, and the main one, promising results through fibromyalgia. With the organization of health systems to adhere to self-applicable or simple inclusive care, Lian Gong seems to successfully meet the purpose of CIPs.

References

1. Prabhakar A, Kaiser JM, Novitch MB, Cornett EM, Urman RD, Kaye AD. The role of complementary and alternative medicine treatments in fibromyalgia: a comprehensive review. *Curr Rheumatol Rep.* 2019;21(5):14.
2. Terzi H, Terzi R, Kale A. The relationship between fibromyalgia and pressure pain threshold in patients with dyspareunia. *Pain Res Manag.* 2015;20(3):137-140.
3. Sarzi-Puttini P, Giorgi V, Marotto D, Atzeni F. Fibromyalgia: an update on clinical characteristics, aetiopathogenesis and treatment. *Nat Rev Rheumatol.* 2020;16(11):645-660.
4. Wolfe F, Clauw DJ, Fitzcharles MA, et al. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res.* 2010;62(5):600-610.
5. Zeng Y, Xie X, Cheng ASK. Qigong or Tai Chi in Cancer Care: an Updated Systematic Review and Meta-analysis. *Curr Oncol Rep.* 2019;21(6):48.
6. Chang P-S, Knobf T, Oh B, Funk M. Physical and psychological health outcomes of Qigong exercise in older adults: A systematic review and meta-analysis. *Am J Chin Med.* 2019;47(02):301-322.
7. Wang CW, Chan CH, Ho RTH, Chan JS, Ng S-M, Chan CLW. Managing stress and anxiety through qigong exercise in healthy adults: a systematic review and meta-analysis of randomized controlled trials. *BMC Complement Altern Med.* 2014;14(1):8.
8. Hwang EY, Chung SY, Cho JH, Song MY, Kim S, Kim JW. Effects of a brief Qigong-based stress reduction program (BQSRP) in a distressed Korean population: a randomized trial. *BMC Complement Altern Med.* 2013;13(1):113.
9. Lin CY, Wei TT, Wang CC, Chen WC, Wang YM, Tsai SY. Acute Physiological and Psychological Effects of Qigong Exercise in Older Practitioners. *Evidence-Based Complement Altern Med.* 2018;2018:1-10.
10. Hartley L, Lee M, Kwong J, et al. Qigong for the primary prevention of cardiovascular disease (Cochrane review) [with consumer summary]. *Cochrane Database Syst Rev.* 2015;11(6):CD010390.
11. Hung LH, Wu CH, Lin BF, Hwang LS. Hyperimmune colostrum alleviates rheumatoid arthritis in a collagen-induced arthritis murine model. *J Dairy Sci.* 2018;101:3778-3787.
12. Lauche R, Cramer H, Häuser W, Dobos G, Langhorst J. A systematic review and meta-analysis of qigong for the fibromyalgia syndrome. *Evidence-Based Complement Altern Med.* 2013;96(45):e8517.
13. Health M of. Ordinance No. 971, of May 03, 2006. Official Federal Gazette. Published 2006. https://bvsmms.saude.gov.br/bvs/saudelegis/gm/2006/prt0971_03_05_2006.html
14. Randow R, Carolina Mendes N, Toyoko Hanashiro Silva L, et al. Lian Gong em 18 terapias como estratégia de promoção da saúde. *Rev Bras em Promoção da Saúde.* 2017;30(4):1-10.
15. Fidalgo J, Zanetini AA, Beraquet B, Bognar MC. Manuais do Lian Gong / Lian Gong Manuals. Prefeitura de Cam

- pinas. Published 2001. http://www.campinas.sp.gov.br/arquivos/gestao-desenvolvimento-pessoas/manual_lian_gong_01.pdf
16. Heymann RE, Paiva ES, Martinez JE, et al. New guidelines for the diagnosis of fibromyalgia. *Rev Bras Reumatol (English Ed)*. 2017;57(S 2):467-476.
 17. Rampakakis E, Ste-Marie PA, Sampalis JS, Karellis A, Shir Y, Fitzcharles MA. Real-life assessment of the validity of patient global impression of change in fibromyalgia. *RMD Open*. 2015;1(1):e000146.
 18. Lauche R, Cramer H, Häuser W, Dobos G, Langhorst J. A systematic overview of reviews for complementary and alternative therapies in the treatment of the fibromyalgia syndrome. *Evidence-based Complement Altern Med*. 2015;2015:610615.
 19. Lorena SB de, Pimentel EA dos S, Fernandes VM, Pedrosa MB, Ranzolin A, Duarte ALBP. Evaluation of pain and quality of life of fibromyalgia patients. *Rev Dor*. 2016;17(1):8-11.
 20. Galvez-s CM, Paso GAR del. Diagnostic criteria for fibromyalgia: Critical review and future perspectives. *J Clin Med*. 2020;9(4):1219.