



REVIEW PAPER

Natalia Leksa ^{1(AEF)}, Anna Pękala ^{2(AEF)}, Piotr Pardak ^{2,3(AEF)}, Rafał Dzedzic ^{4(AEF)},
Dorota Bartusik-Aebisher ^{5(BCDEF)}, Rafał Filip ^{2,3(AEF)}, Sabina Galiniak ^{5(AEF)},
David Aebisher ^{6(ADEF)}

The potential of intracellular ¹³C MR spectroscopy to study the absolute configuration of endogenous and polarized alanine

¹ Department of Human Anatomy, Medical College of Rzeszów University, Rzeszów, Poland

² Department of Gastroenterology with IBD Unit, Clinical Hospital No. 2, Rzeszów, Poland

³ Department of Gastroenterology, Medical College of Rzeszów University, Rzeszów, Poland

⁴ Department of Neurosurgery, Hospital MSWiA, Rzeszów, Poland

⁵ Department of Biochemistry and General Chemistry, Medical College of Rzeszów University, Rzeszów, Poland

⁶ Department of Photomedicine and Physical Chemistry, Medical College of Rzeszów University, Rzeszów, Poland

ABSTRACT

Introduction. Quantitative and accurate monitoring of tumor makes hyperpolarized carbon (¹³C) Magnetic Resonance Imaging and Spectroscopy (MRI/S) a powerful tool for in vivo metabolic and structural study. Moreover, the studies of the properties and functions in tumor tissue of the compounds of carbon (C) that are organic, are fundamental to tumor biochemistry.

Aim. To review ¹³C MR spectroscopy to study the absolute configuration of endogenous and polarized alanine

Material and methods. An analysis of literature regarding ¹³C MR spectroscopy of polarized alanine.

Results. Current evidence suggests that the determination of absolute configurations of amino acids play significant role in physiological mechanisms during tumor growth and treatment.

Conclusions. Nearly 50% nuclear polarization for ¹³C can be achieved in various organic molecules when Dynamic Nuclear Polarization DNP is performed in a strong magnetic field and at cryogenic temperatures.

Keywords. ¹³C NMR, dynamic nuclear polarization, hyperpolarized carbon

Introduction

From all types of spectroscopy, Magnetic Resonance Spectroscopy (MRS) can provide a useful analytical tool to determine the absolute configuration (L,D) of some metabolites in biomedical assays.¹⁻⁵ All enzyme cata-

lyzed reactions in living matter are stereospecific and provide product in enantiomeric forms. By definition, the enantiomeric forms are mirror images of each other and their high-resolution spectra recorded in achiral media are identical. However, two enantiomers will

Corresponding author: Dorota Bartusik-Aebisher, e-mail: dbartusik-aebisher@ur.edu.pl

Participation of co-authors: A – Author of the concept and objectives of paper; B – collection of data; C – implementation of research; D – elaborate, analysis and interpretation of data; E – statistical analysis; F – preparation of a manuscript; G – working out the literature; H – obtaining funds

Received: 6.11.2020 | Accepted: 13.01.2020

Publication date: March 2020

lose the same chemical potential and consequently will have differ MRS spectrum, while interacting with a further chiral entity.⁶⁻⁹ Moreover, observation of enantiomeric composition of samples, based on MRS in the presence of the chiral shift reagent can induce precise determination of the ratios of both enantiomers and induce its separation. The chiral entity cannot only separate two forms but it also has influence on the sensitivity of recorded spectra by establishing labile diastereomeric interactions with enantiomers.¹⁰⁻¹⁵ The split of the resonances of the two forms can be exploited to assess quantitatively the enantiomeric composition of a living matter. It is obvious that stereochemical considerations of the enantiomeric forms are of high importance for understanding biochemical and physiological mechanisms in cancer tissue. Determination of absolute structure of metabolites, such as amino acids, might be a sufficient way to early detection of metabolic disorders that has implications for diagnosis and treatment. In particular, in order to target a biological receptor, the specificity and chirality of the recognition site must be correct. When such peptides are composed of L-amino acids, they are rapidly degraded *in vivo* by naturally occurring proteases but when composed of D-amino acids, the peptide is resistant to degradation, giving the possibility for therapeutic use.¹⁶⁻²⁰

Molecular asymmetry

Molecular asymmetry is fundamental for intracellular processes of chiral components in both healthy and cancer tissue. A molecule is chiral when it cannot be superimposable on its mirror image with the chiral object and its mirror image cannot be referred to as enantiomers. Almost all intracellular reactions are catalyzed by enzyme and usually provide two enantiomeric products with the absolute configuration of right-handed (D-) configuration and left-handed (L-) configuration. Recent studies have shown that both, D- and L- amino acids (AA) are present in animal and human body in high concentrations as building blocks of proteins, intermediates in metabolism and fulfill specific physiological functions. However, the structure and symmetry of organic molecules also play crucial role in many biochemical processes. One of the reasons for the increasing interest in the separation and quantification of both AA configurations lies in clinical diagnostics. The pros and cons of existing methods for enantioselective analysis of AA and their applications to biological samples are recent trends in the field but there are several research questions.²¹⁻²⁸

^{13}C MRS

In particular, breast cancer cells show varied concentrations of AA using mostly chromatographic techniques; however their chirality was not studied yet. Therefore, the detection of D- and L- AA is of particular interest for

our study since the determination of absolute configurations play significant role in physiological mechanisms during tumor growth and treatment. It is generally held that breast cancer cells express strong glycolytic activity and release large amounts of lactate, which can be produced or consumed with regard to the oxygenation of cells. D- and L-alanine was already found in gland and epithelial cells of mammals, but is typically quantified as the sum of their D- and L-enantiomers.²⁹⁻³³ Moreover, alanine is known to be the predominant amino acid for the growth of breast cancer cells. Because the intracellular functions of alanine such as cell regeneration by L- form and damage by D- form are associated with absolute configurations, their ratio can be useful in diagnostic and therapeutic processes. We focused on the detection of absolute configuration of intracellular alanine *ex vivo* using carbon-13 magnetic resonance spectroscopy (^{13}C MRS). The chemical shift range for ^{13}C (~250 ppm) is much larger than that for proton (~15 ppm) and the T_1 relaxation time of ^{13}C in small molecules is much longer than that of ^1H . However, in nature, about 1.11% of naturally occurring C is ^{13}C isotope, which is magnetically active and can be applied to probe molecular structures that correspond to physiological changes in tumor tissues. More than 98.89% of naturally occurring C is ^{12}C with no MR signal.³⁵⁻⁴¹ Several techniques have been used to overcome the ^{13}C MRS limitation by means of enhancing the polarization of nuclear spins. One of the techniques is proton decoupling, which eliminates the coupling of ^1H with ^{13}C by irradiating the entire ^1H resonance absorption range and consequently collapsing ^{13}C resonances to singlets. Although the MR signal of ^{13}C measured intracellularly corresponds to metabolic changes, signal intensity (SI) is too low to be relevant for quantitative and longitudinal study. However, the visualization of ^{13}C nuclei concentration may result in images and spectra with high signal to noise ratio (SNR) due to the hyperpolarization process. Nearly 100% nuclear polarization for ^1H and 50% for ^{13}C can be achieved in various organic molecules when Dynamic Nuclear Polarization (DNP) is performed in a strong magnetic field and at cryogenic temperatures. Replacing the ^{12}C (98.9% natural abundance) isotope with the ^{13}C isotope at a specific carbon or carbons in a metabolic substrate does not affect the substrate's biochemistry. With ^{13}C MRS, the body tissues are virtually invisible, and only regions where the hyperpolarized ^{13}C -labeled substance is present will appear in the generated images.³⁶⁻³⁹ In nature, C is abundant in all forms of life and all dead organic materials. Although the MR signal of ^{13}C *in vivo* corresponds to metabolic changes, signal intensity (SI) of naturally occurring ^{13}C is too low to be relevant for quantitative and longitudinal study. However, the visualization of ^{13}C nuclei concentration may result in images and spectra with high signal to noise ratio (SNR) due to the hyperpolar-

ization process. Hyperpolarization does not change any chemical or physical properties of the substances, however it allows acquisition of ^{13}C images and spectra in a relevant time frame. Hence, hyperpolarized ^{13}C MRI/S can directly inform about the biochemical tissue composition and chemical structure by generating frequency and spatial distribution of hyperpolarized atoms. In general, most suitable ^{13}C compounds for MR are small molecules (molecular weight $\sim 120\text{ gmol}^{-1}$) with a possibility to obtain information about molecular behavior *in vivo* due to rapid uptake in tissue. We review the applications of ^{13}C hyperpolarized techniques such as dynamic-nuclear-polarization (DNP) to monitor tumor targeting giving an overview of rapid ^{13}C MRI/S sequences used which is followed by the hardware enhancement such as coil design.²⁹⁻³³ Hyperpolarization does not change any chemical or physical properties of the substances; however it allows acquisition of ^{13}C images and spectra within a relevant time frame. Hence, hyperpolarized ^{13}C MRI/S can directly inform about the biochemical tissue composition and chemical structure by generating frequency and spatial distribution of hyperpolarized atoms. In general, most suitable ^{13}C compounds for MR are small molecules (molecular weight $\sim 120\text{ gmol}^{-1}$) with a possibility to obtain information about molecular behavior *in vivo* due to rapid uptake in tissue. Prior to ^{13}C MRS *ex vivo* experiments, can be use presently available ^{13}C D-, L- alanine and ^{13}C L- Alanine to confirm ^{13}C resonance assignment in water solution phantoms. In these studies, we propose the utilization of at least three chiral lanthanide complexes such as (1) lanthanum(III)-tris (6,6,7,7,8,8,8-heptafluoro-2,2-dimethyl-3,5-octanedionate, $\text{C}_{30}\text{H}_{30}\text{F}_{21}\text{LaO}_6$, (2) europium tris[3-(heptafluoropropylhydroxymethylene)-(-)-camphorate, $\text{C}_{42}\text{H}_{42}\text{EuF}_{21}\text{O}_6$ and (3) ytterbium(III)-tris[3-(heptafluoropropylhydroxymethylene)-l-camphorate, $\text{C}_{42}\text{H}_{42}\text{F}_{21}\text{O}_6\text{Yb}$, as a probe for differentiation of the enantiomeric chirality on the racemic D- and L- spectra and enantiomeric (D or L) spectra of amino acids solution.³³⁻³⁶ Proposed chemical shift agents are highly water-soluble and suitable for study in the living matter. The chiral entity (1-3) cannot only separate two absolute forms of D- and L- alanine but it also has influence on the sensitivity of the recorded spectra by establishing labile diastereomeric interactions with enantiomers. The split of resonances of the two forms can be exploited to assess quantitatively the enantiomeric composition of the living matter.³⁷⁻⁴¹

Conclusion

Due to the beneficial impact of intracellular ^{13}C MR spectroscopy to study the absolute configuration of endogenous and polarized alanine in medicine there is a growing interest in analytical identification and quantification *in vitro* and *in vivo*.

Acknowledgements

Dorota Bartusik-Aebisher acknowledges support from the National Center of Science NCN (New drug delivery systems-MRI study, Grant OPUS-13 number 2017/25/B/ST4/02481).

References

- Brandes AH, Ward CS, Ronen SM. 17-allylamino-17-demethoxygeldanamycin treatment results in a magnetic resonance spectroscopy-detectable elevation in choline-containing metabolites associated with increased expression of choline transporter SLC44A1 and phospholipase A2. *Breast Cancer Res.* 2010;12(5):R84.
- Jagannathan NR, Sharma U. Breast Tissue Metabolism by Magnetic Resonance Spectroscopy. *Metabolites.* 2017;7(2):23-27.
- Maher EA, Marin-Valencia I, Bachoo RM, et al. Metabolism of [U- ^{13}C]glucose in human brain tumors *in vivo*. *NMR Biomed.* 2012;25(11):1234-44.
- Grinde MT, Moestue SA, Borgan E, Risa Ø, Engebraaten O, Gribbestad IS. ^{13}C high-resolution-magic angle spinning MRS reveals differences in glucose metabolism between two breast cancer xenograft models with different gene expression patterns. *NMR Biomed.* 2011;24(10):1243-1252.
- Wang W, Yi F, Ni Y, Zhao Z, Jin X, Tang, Y. Parity violation of electroweak force in phase transitions of single crystals of mD- and L- alanine and valine. *J. Biol. Phys.* 2000;26:51-65.
- Wang WQ, Min W, Bai F, Sun L, Yi F, Wang ZM, Yan CH, Ni YM, Zhao ZX. Temperature dependent magnetic susceptibilities study on parity violating phase transitions of D- and L-alanine crystals. *Tetrahedron Asymm.* 2002;13:2427-2432.
- Wang WQ, Min W, Liang Z, Wang LY, Chen L, Deng F. NMR and parity violation: low temperature dependence in ^1H CRAMPS and ^{13}C CP/MAS ssNMR spectra of alanine enantiomer. *Biophys. Chem.* 2003;103:289-298.
- Sullivan R, Pyda M, Pak J, et al. Search for electroweak interactions in amino acid crystals. II The Salam hypothesis. *J Phys Chem A.* 2003;107:6674-6680.
- Smith GG, Reddy GV. Effect of side chain on the racemisation of amino acids in aqueous solution. *J Org Chem.* 1989;54:4529-4535.
- Ahmed AF, Teng WT, Huang CY, Dai CF, Hwang TL, Sheu JH. Anti-Inflammatory Lobane and Prenyleudesmane Diterpenoids from the Soft Coral *Lobophytum varium*. *Mar Drugs.* 2017;15(10): E300.
- Rogge B, Itagaki Y, Fishkin N, et al. Retinoylserine and retinoylalanine, natural products of the moth *Trichoplusia ni*. *J Nat Prod.* 2005;68(10):1536-1540.
- Moon CM, Kim YH, Ahn YK, Jeong MH, Jeong GW. Metabolic alterations in acute myocardial ischemia-reperfusion injury and necrosis using *in vivo* hyperpolarized [1- ^{13}C] pyruvate MR spectroscopy. *Sci Rep.* 2019;9(1):18427.

13. Can E, Mishkovsky M, Yoshihara HAI, et al. Noninvasive rapid detection of metabolic adaptation in activated human T lymphocytes by hyperpolarized ¹³C magnetic resonance. *Sci Rep.* 2020; 13;10(1):200.
14. Glunde K, Jie C, Bhujwalla ZM. Select item 20641983ses of the aberrant choline phospholipid metabolism in breast cancer. *Cancer Res.* 2004;64(12):4270-4276.
15. Rohm M, Savic D, Ball V, et al. Cardiac Dysfunction and Metabolic Inflexibility in a Mouse Model of Diabetes Without Dyslipidemia. *Diabetes.* 2018;67(6):1057-1067.
16. Eills J, Cavallari E, Carrera C, Budker D, Aime S, Reineri F. Real-Time Nuclear Magnetic Resonance Detection of Fumarate Activity Using Parahydrogen-Hyperpolarized [1-¹³C]Fumarate. *J Am Chem Soc.* 2019;141(51):20209-20214.
17. Glöggler S, Rizzitelli S, Pinaud N, et al. In vivo online magnetic resonance quantification of absolute metabolite concentrations in microdialysate. *Sci Rep.* 2016;4,6:36080.
18. Ward CS, Eriksson P, Izquierdo-Garcia JL, Brandes AH, Ronen SM. HDAC inhibition induces increased choline uptake and elevated phosphocholine levels in MCF7 breast cancer cells. *PLoS One.* 2013;8(4):e62610.
19. Darpolor MM, Kaplan DE, Pedersen PL, Glickson JD. Human Hepatocellular Carcinoma Metabolism: Imaging by Hyperpolarized ¹³C Magnetic Resonance Spectroscopy. *J Liver Disease Transplant.* 2012;1(1). doi: 10.4172/2325-9612.1000101.
20. Kim S, Martinez-Santesteban F, Scholl TJ. Measuring the Spin-Lattice Relaxation Magnetic Field Dependence of Hyperpolarized [1-¹³C]pyruvate. *J Vis Exp.* 2019;(151).
21. Lee CY, Soliman H, Geraghty BJ, et al. Lactate topography of the human brain using hyperpolarized ¹³C-MRI. *Neuroimage.* 2020;204:116202.
22. van de Vena KCC, van der Graafa M, Tack CJJ, et al. Optimized [1-¹³C]glucose infusion protocol for ¹³C magnetic resonance spectroscopy at 3 T of human brain glucose metabolism under euglycemic and hypoglycemic conditions. *J Neur Methods.* 2010;186:68-71.
23. Yanga J, Johnsona C, Shena J. Detection of reduced GABA synthesis following inhibition of GABA transaminase using in vivo magnetic resonance signal of [¹³C]GABA C1. *Journal of Neuroscience Methods.* 2009;182:236-243.
24. Schroeder MA, Swietach P, Atherton HJ, et al. Measuring intracellular pH in the heart using hyperpolarized carbon dioxide and bicarbonate: a ¹³C and ³¹P magnetic resonance spectroscopy study. *Cardiovascular Research.* 2010;86:82-91.
25. Mayer D, Yen YF, Tropp J, Pfefferbaum A, Hurd RE, Spielman DM. Application of subsecond spiral chemical shift imaging to real-time multislice metabolic imaging of the rat in vivo after injection of hyperpolarized ¹³C1-pyruvate. *Magn Reson Med.* 2009;62(3):557-564.
26. Keshari KR, Wilson DM, Chen AP, et al. Hyperpolarized [2-¹³C]-fructose: a hemiketal DNP substrate for in vivo metabolic imaging. *J Am Chem Soc.* 2009;131(48):17591-17596.
27. Jamin Y, Gabellieri C, Smyth L, et al. Hyperpolarized (¹³C) magnetic resonance detection of carboxypeptidase G2 activity. *Magn Reson Med.* 2009;62(5):1300-1304.
28. Park I, Larson PE, Zierhut ML, et al. Hyperpolarized ¹³C magnetic resonance metabolic imaging: application to brain tumors. *Neuro Oncol.* 2010;12(2):133-144.
29. Bhattacharya P, Harris K, Lin AP, et al. Ultra-fast three dimensional imaging of hyperpolarized ¹³C in vivo. *MAGMA.* 2005;18(5):245-256.
30. Aime S, Dastrù W, Gobetto R, Santelia D, Viale A. Agents for polarization enhancement in MRI. *Handb Exp Pharmacol.* 2008;(185 Pt 1):247-272.
31. Schröder L, Lowery TJ, Hilty C, Wemmer DE, Pines A. Molecular imaging using a targeted magnetic resonance hyperpolarized biosensor. *Science.* 2006;314(5798):446-449.
32. Bhattacharya P, Chekmenev EY, Perman WH, et al. Towards hyperpolarized (¹³C)-succinate imaging of brain cancer. *J Magn Reson.* 2007;186(1):150-155.
33. Ross BD, Bhattacharya P, Wagner S, Tran T, Sailasuta N. Hyperpolarized MR imaging: neurologic applications of hyperpolarized metabolism. *AJNR Am J Neuroradiol.* 2010;31(1):24-33.
34. Golman K, Petersson JS, Magnusson P, et al. Cardiac metabolism measured noninvasively by hyperpolarized ¹³C MRI. *Magn Reson Med.* 2008;59(5):1005-1013.
35. Hancu I, Wood SJ, Piel J, et al. Three-frequency RF coil designed for optimized imaging of hyperpolarized, ¹³C-labeled compounds. *Magn Reson Med.* 2008;60(4):928-933.
36. Mayer D, Yen YF, Levin YS, et al. In vivo application of sub-second spiral chemical shift imaging (CSI) to hyperpolarized ¹³C metabolic imaging: comparison with phase-encoded CSI. *J Magn Reson.* 2010;204(2):340-345.
37. Lingwood MD, Han S. Dynamic nuclear polarization of ¹³C in aqueous solutions under ambient conditions. *J Magn Reson.* 2009;201(2):137-145.
38. Marjańska M, Iltis I, Shestov AA, et al. In Vivo ¹³C Spectroscopy in the Rat Brain using Hyperpolarized [1-¹³C]pyruvate and [2-¹³C]pyruvate. *J Magn Reson.* 2010;206(2):210-218.
39. Hu S, Chen AP, Zierhut ML, et al. In vivo carbon-13 dynamic MRS and MRSI of normal and fasted rat liver with hyperpolarized ¹³C-pyruvate. *Mol Imaging Biol.* 2009;11(6):399-407.
40. Chen AP, Leung K, Lam W, Hurd RE, Vigneron DB, Cunningham CH. Design of spectral-spatial outer volume suppression RF pulses for tissue specific metabolic characterization with hyperpolarized ¹³C pyruvate. *J Magn Reson.* 2009;200(2):344-348.
41. Ferdia A, Gallagher, Mikko I, Kettunen, Sam E. ¹³C MR Spectroscopy Measurements of Glutaminase Activity in Human Hepatocellular Carcinoma Cells Using Hyperpolarized ¹³C-Labeled Glutamine. *Mag Res Med.* 2008; 60:253-257.